

What type of graph is best for my data?



Graphs are the best way to display data when overall patterns are more important than individual data values. The shape of the graph reveals trends, differences, exceptions, anomalies and other relationships that might be difficult to detect in a table.

Always choose the correct type of graph for your data so that these relationships are clear. Usually, the simplest graph will be best. Readers will engage with, and be persuaded by, the messages if your data are presented in a way that matches their intuitive understanding of data relationships. For example, left-to-right lines are an intuitive way to represent measurements over time. Some relationships can be shown by more than 1 type of graph. In this case, you should use the graph type that is most familiar to your readership.





The next few sections show various data relationships and recommended graph types for each relationship.¹

Change over time

Shows how data values for 1 or more measures change over time (e.g. population-adjusted breast cancer diagnoses recorded in Australia every year, for the past 20 years).




Recommended graph type	Features and notes
 <p>Line graph (for large time series)</p>	<p>Use to highlight trends or patterns in a measure over many time points</p> <p>Use for datasets that include data for more than about 8 time points</p> <p>Lines connect consecutive data values</p> <p>Lines always go left to right, with time intervals on the x axis increasing from left to right, and the measurement variable plotted against the y axis</p> <p>Only connect consecutive values – intervals with missing data must be shown as breaks in the line</p>
 <p>Vertical bar graph (for small time series)</p>	<p>Use for time-series data with a small number of time points (around 8 or fewer)</p> <p>Use to emphasise specific data values, rather than an overall pattern or trend</p>

¹ Examples are adapted from Few S (2012). *Show me the numbers: designing tables and graphs to enlighten*, 2nd edn, Analytics Press, Burlingame, California; and Evergreen SDH (2017). *Effective data visualization: the right chart for the right data*, SAGE Publications, Thousand Oaks, California.

Recommended graph type	Features and notes
 <p data-bbox="134 584 264 618">Dot graph</p>	<p data-bbox="531 282 1098 315">Dots represent data values at each time point</p> <p data-bbox="531 327 1458 398">Can be mistaken for scatter plots – consider using a vertical bar graph or a line graph instead</p>
 <p data-bbox="134 936 339 969">Dumbbell graph</p>	<p data-bbox="531 633 1426 705">Connected dots represent 2 time points of data (e.g. pre- and post-test) for several groups</p> <p data-bbox="531 716 1406 788">Use with care; this graph type is less common and will be unfamiliar to many readers</p>
 <p data-bbox="134 1288 448 1321">Vertical bar cluster graph</p>	<p data-bbox="531 985 1347 1019">Can be used to show data for several groups at 2 or 3 time points</p> <p data-bbox="531 1030 1415 1102">Can be difficult for readers to compare differences across many groups because of distance between groups on x axis</p>
 <p data-bbox="134 1639 442 1711">Trellis of single-category graphs</p>	<p data-bbox="531 1337 1442 1449">Consider using a trellis (panel) graph with a small graph for each group if you have many groups or categories of data and they would appear cluttered if they were all presented on the same line or bar graph</p> <p data-bbox="531 1460 1430 1494">Keep the axis ranges consistent across all graphs to enable comparisons</p>




Comparing groups or categories

Compares data values across independent items, groups or categories (e.g. unemployment rates for each Australian state and territory).

Recommended graph type	Features and notes
	<p>If categories have no inherent order, can order bars by size of data values to emphasise differences</p>
<p>Horizontal bar graph</p>	
	<p>Use clustered bars for subcategories of groups, but limit clusters to 3 or 4 subcategories to enable comparisons across groups</p>
<p>Horizontal or vertical bar cluster graph</p>	
	<p>Connected dots represent 2 related data points on a common scale Readers can easily judge distance between dots, and differences in this distance across groups Use with care; this graph type is less common than the others and will be unfamiliar to many readers</p>
<p>Dumbbell graph</p>	





Single frequency or distribution data

Shows how possible values are distributed over the range of a measure (e.g. range of blood pressure measurements for people aged 60 to 70 years).

Recommended graph type	Features and notes
 <p data-bbox="134 730 526 801">Histogram (for measures with a few possible discrete values)</p>	<p data-bbox="587 434 1457 506">Use a histogram to show frequencies or count values across the range of a measure with few intervals</p> <p data-bbox="587 517 1457 589">Use as an alternative to a frequency polygon when individual data values must be emphasised</p>
 <p data-bbox="134 1124 552 1196">Frequency polygon (for measures with many possible values)</p>	<p data-bbox="587 822 1457 893">Use to show frequency or count values across the range of a measure with many intervals</p> <p data-bbox="587 904 1139 936">Use to emphasise the shape of a distribution</p>
 <p data-bbox="134 1512 252 1543">Strip plot</p>	<p data-bbox="587 1209 1385 1240">Use to show the distribution of a measure for a small population</p> <p data-bbox="587 1256 1426 1328">If many measurements have the same value, these points should be stacked or shown in a denser tone than nonrepeated points</p> <p data-bbox="587 1339 1433 1447">Use with care; this graph type is less common than some others and will be unfamiliar to many readers. Consider plotting a simple histogram instead</p>
 <p data-bbox="134 1863 517 1895">Box plot (horizontal or vertical)</p>	<p data-bbox="587 1561 1407 1632">Use to show a summary of a measure's distribution, rather than all individual data values</p> <p data-bbox="587 1644 1401 1751">Use with care; this graph type is less common than some others and will be unfamiliar to many readers. Consider plotting a simple histogram instead</p>





Part to whole (or proportions of a whole)

Shows how data values relate to, compare with or make up a total measure at 1 or more times, places and so on (e.g. proportion of Australia's total primary energy supply attributable to each major fuel type).

Recommended graph type	Features and notes
 <p data-bbox="134 730 395 770">Horizontal bar graph</p>	<p data-bbox="587 434 1420 501">Use to show the value (i.e. percentage or proportion of an absolute total) of each part for a single population</p> <p data-bbox="587 515 1340 582">This type of data is often shown as a pie graph, which is NOT recommended for displaying scientific data</p>
 <p data-bbox="134 1079 497 1120">Horizontal stacked bar graph</p>	<p data-bbox="587 784 1452 851">Use to show proportions of a total measure for several populations or groups</p> <p data-bbox="587 864 1398 931">Total(s) must add to 100% if parts are percentages, or to the total absolute value for other measures</p>
 <p data-bbox="134 1438 459 1469">Vertical stacked bar graph</p>	<p data-bbox="587 1133 1410 1200">Use to show proportions of a total measure for several times, populations or groups, for about 8 or fewer populations or groups</p> <p data-bbox="587 1214 1398 1281">These groups may be the same group, measured at various times</p> <p data-bbox="587 1258 1385 1294">Use to emphasise changes in the relative sizes of parts over time</p>
 <p data-bbox="134 1787 376 1823">Stacked area graph</p>	<p data-bbox="587 1482 1452 1550">Use to show proportions of a total measure over time, for around 8 or more time points</p> <p data-bbox="587 1572 1385 1608">Use to emphasise changes in the relative sizes of parts over time</p>


Distribution of the same measure across many time points or categories

Shows how frequency or count values are distributed across the range of a measure, possibly for more than 1 population (e.g. range of blood pressure measurements for people aged 60 to 70 years with 5 different medical conditions).

Recommended graph type	Features and notes
 <p data-bbox="134 770 541 837">Line graph with upper and lower bounds</p>	<p data-bbox="584 472 1410 539">Use to show the distribution of a measure as it varies with time for many time points</p> <p data-bbox="584 555 1457 622">Summary values (e.g. median or mean values) for the measure at each time point are connected to form a line</p> <p data-bbox="584 638 1442 779">The upper confidence limits at each time are connected to form a (typically invisible) line above the data, and the lower confident limits are connected to form a line below the data, and the area between these lines is shaded</p> <p data-bbox="584 795 1406 902">Upper and lower bounds may be unfamiliar to readers – consider whether their inclusion adds meaning and whether this outweighs potential misunderstandings</p>
 <p data-bbox="134 1218 252 1252">Strip plot</p>	<p data-bbox="584 920 1337 954">Distributions are plotted side by side against the same y axis</p> <p data-bbox="584 969 1145 1003">Use white space to separate the distributions</p> <p data-bbox="584 1019 1457 1086">Use to show the distribution of a measure for a small population for a few categories or times on the x axis</p> <p data-bbox="584 1102 1442 1243">If several measurements have the same value for a given distribution, these points should be stacked (for example, moved sideways by differing amounts) or shown in a denser tone than nonrepeated points</p>
 <p data-bbox="134 1572 341 1606">Vertical box plot</p>	<p data-bbox="584 1274 1353 1341">Use to summarise multiple distributions of the same measure, perhaps over time or across categories</p> <p data-bbox="584 1357 1442 1498">May be unfamiliar to readers – consider plotting summary values (e.g. medians) as a bar graph for multiple groups or populations, or a line graph with or without upper and lower bounds for multiple distributions over time</p>
 <p data-bbox="134 1924 533 1957">Trellis of single-category graphs</p>	<p data-bbox="584 1628 1442 1769">Consider using a trellis (panel) graph with a small graph for each group if you have many groups or categories of data and they would appear cluttered if they were all presented on the same line or bar graph</p> <p data-bbox="584 1785 1426 1852">Keep the axis ranges consistent across all graphs to enable accurate comparisons</p>



Deviation

Shows the difference between data values and a baseline, target or threshold (e.g. differences between actual rainfall and predicted or previous-year rainfall for each month of a year).

Recommended graph type	Features and notes
 <p>Vertical bar graph with baseline</p>	<p>Use when the goal is to highlight deviations of measurements from some meaningful baseline or reference</p> <p>Bars (i.e. data values) above the reference or x axis indicate positive differences from the baseline; bars below indicate negative differences</p> <p>The y axis can measure absolute differences or percentage change between data values and the reference</p>
 <p>Line graph with baseline</p>	<p>Use to show differences from a baseline or reference over time when the dataset includes data for more than about 8 time points</p> <p>Use to highlight trends in a measure over many time points</p> <p>Lines are connected, consecutive data values</p> <p>Lines always go left to right, with time on the x axis increasing from left to right, and the measurement variable plotted against the y axis</p> <p>Only connect consecutive values – missing data must be shown by breaks in the line</p>

Correlated measures

Shows an association between 2 measures or variables (e.g. children's ages and heights).

Recommended graph type	Features and notes
 <p>Scatter plot</p>	<p>Each dot or data point represents a subject's measurement on x-axis and y-axis variables</p> <p>Use to show that data points form a meaningful shape that indicates the type (or lack) of association between 2 variables</p> <p>Consider including a trend line to highlight the type and strength of association</p> <p>Some readers may struggle to interpret scatter plots – consider whether side-by-side or stacked horizontal bar graphs would better communicate the association</p>
 <p>Side-by-side horizontal bar graph</p>	<p>Use to show an association between 2 measures when scatter plots are unfamiliar to readers</p> <p>Most effective for showing linear associations</p> <p>Two aligned bar graphs display each subject's measurements on the first and second measures</p> <p>Order the bars by size on 1 of the graphs to emphasise the association between the 2 measures</p>